

| Led | Sortskode | Sortsnavn | Hybrid / Linie | Udbytte | | | | | Standardkvalitet | | | | | Dyrkningsegenskaber | | | | Kvalitetsegenskaber | | | | | | | Oliekvalitet, pct. (%) | | | | | |
|-----|-----------|------------|----------------|-----------------------------------|--------|-------------|------|-----|------------------|--------|-------------|------|-----|---------------------|------------------|----------------------------|----------------------|---------------------|----------------------|----------------|---------------------|------------------|-------------|-----------------|-----------------------------|------------------|-------------------|---------------------|--------------------|--|
| | | | | hkg/ha korrigeret til 91% tørstof | | | | | hkg/ha | | | | | | | | | Frøkvalitet | | | | | | | | | | | | |
| | | | | Abildgård | Holeby | Koldkærgård | Gns. | FHT | Abildgård | Holeby | Koldkærgård | Gns. | FHT | Blomstring, dato | Plante højde, cm | Afgroede højde v. høst, cm | Lejesæd, skala: 0-10 | Frøvægt, mg pr. frø | Proteinindhold, pct. | Protein, kg/ha | Proteinindhold, FHT | Olieindhold, pct | Olie, kg/ha | Olie, kg/ha FHT | Glucosinolatindhold, µmol/g | Oliesyre, C 18:1 | Linolsyre, C 18:2 | Linolensyre, C 18:3 | Eruceasyre, C 22:1 | |
| | | Amtal fs. | | | | | 3 | 3 | | | | 3 | 3 | 2 | 1 | 1 | 1 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 2 | 1 | 1 | 1 | 1 | |
| 1 | 9018 | Blanding | DH | 50.0 | 51.4 | 41.4 | 47.6 | 100 | 53.1 | 54.5 | 43.8 | 50.5 | 100 | 0305 | 172 | 170 | 0.0 | 4.0 | 18.9 | 817 | 100 | 49.2 | 2132 | 100 | | | | | | |
| 2 | 30641 | PX128 | DH | 35.2 | 41.0 | 37.0 | 37.7 | 79 | 38.3 | 44.5 | 39.5 | 40.8 | 81 | 0605 | 123 | 130 | 0.0 | 4.5 | 18.5 | 636 | 78 | 51.0 | 1751 | 82 | | | | | | |
| 3 | 33865 | X18WX010C | DH | 30.2 | 41.8 | 37.7 | 36.6 | 77 | 32.0 | 44.8 | 40.2 | 39.0 | 77 | 0605 | 137 | 138 | 0.0 | 4.1 | 19.2 | 639 | 78 | 49.7 | 1656 | 78 | 12.5 | 61.7 | 17.0 | 9.6 | 0.0 | |
| 4 | 33867 | X18WX291C | DH | 35.2 | 41.2 | 33.2 | 36.5 | 77 | 37.6 | 43.8 | 35.0 | 38.8 | 77 | 0705 | 143 | 150 | 0.0 | 3.9 | 19.7 | 656 | 80 | 49.4 | 1642 | 77 | 10.2 | 60.4 | 19.4 | 9.0 | 0.0 | |
| 5 | 33868 | X18WX292C | DH | 35.3 | 40.3 | 37.6 | 37.7 | 79 | 37.7 | 43.1 | 40.1 | 40.3 | 80 | 0405 | 128 | 133 | 0.0 | 4.5 | 19.0 | 653 | 80 | 49.9 | 1714 | 80 | 11.4 | 63.3 | 15.5 | 9.0 | 0.1 | |
| 6 | 33907 | CWH518 | DH | 43.3 | 41.4 | 37.6 | 40.8 | 86 | 46.2 | 44.2 | 39.9 | 43.4 | 86 | 0905 | 142 | 151 | 0.0 | 4.6 | 19.3 | 715 | 88 | 49.7 | 1845 | 87 | 13.1 | 57.6 | 20.6 | 9.8 | 0.1 | |
| 7 | 34625 | X19WX475C | DH | 35.3 | 39.8 | 36.7 | 37.3 | 78 | 37.8 | 42.3 | 38.7 | 39.6 | 78 | 0705 | 135 | 136 | 0.0 | 3.7 | 18.9 | 642 | 79 | 49.4 | 1675 | 79 | 7.5 | 59.6 | 20.4 | 8.5 | 0.1 | |
| 8 | 34618 | X19WT278C | H | 48.3 | 54.7 | 41.9 | 48.3 | 101 | 52.2 | 58.7 | 43.8 | 51.6 | 102 | 0405 | 185 | 191 | 0.0 | 4.2 | 19.4 | 852 | 104 | 49.8 | 2194 | 103 | 7.8 | 60.5 | 18.8 | 9.2 | 0.1 | |
| 9 | 34627 | X19WX479C | DH | 37.6 | 42.9 | 34.9 | 38.5 | 81 | 39.5 | 44.6 | 37.0 | 40.4 | 80 | 0505 | 133 | 135 | 0.0 | 4.3 | 19.9 | 696 | 85 | 48.4 | 1691 | 79 | 8.6 | 58.6 | 19.7 | 9.9 | 0.1 | |
| | | LSD 0.05 | | 4.5 | 6.1 | 2.7 | 2.6 | 5 | | | | | | | | | | | | | | | | | | | | | | |
| | | GNS UDBYTT | | 38.9 | 43.8 | 37.6 | | | | | | | | | | | | | | | | | | | | | | | | |

Udbytte i hkg/ha korrigeret til 91 % tørstof

| Abildgård | Holeby | Koldkærgård | Gns. | Rækkefølge |
|---------------|---------------|---------------|---------------|------------|
| 50.0 Blanding | 106 X19WT278C | 101 X19WT278C | 101 X19WT278C | 1 |
| 97 X19WT278C | 51.4 Blanding | 41.4 Blanding | 47.6 Blanding | 2 |
| 87 CWH518 | 83 X19WX479C | 91 X18WX010C | 86 CWH518 | 3 |
| 75 X19WX479C | 81 X18WX010C | 91 X18WX292C | 81 X19WX479C | 4 |
| 71 X18WX292C | 81 CWH518 | 91 CWH518 | 79 PX128 | 5 |
| 71 X19WX475C | 80 X18WX291C | 89 PX128 | 79 X18WX292C | 6 |
| 70 PX128 | 80 PX128 | 89 X19WX475C | 78 X19WX475C | 7 |
| 70 X18WX291C | 78 X18WX292C | 84 X19WX479C | 77 X18WX010C | 8 |
| 60 X18WX010C | 77 X19WX475C | 80 X18WX291C | 77 X18WX291C | 9 |
| 11 LSD 0.05 | 15 LSD 0.05 | 7 LSD 0.05 | 5 LSD 0.05 | 10 |

Translations

| | | | |
|------------------------------------|---|------------------|-------------------------|
| Afgrødehøjde | <i>Crop height</i> | Stivelsesindhold | <i>Starch content</i> |
| Blomstring | <i>Flowering</i> | Strålængde, cm | <i>Straw length</i> |
| Blødgøring | <i>Softening</i> | Udbytte | <i>Yield</i> |
| Brunrust | <i>Brown rust (Puccinia recondita)</i> | Vandoptagelse | <i>Water absorption</i> |
| Brødhøjde | <i>Bread height</i> | | |
| Brødvolumen | <i>Bread volume</i> | | |
| Bygrust | <i>Barley Rust (Puccinia hordei)</i> | | |
| Dyrkningsegenskaber | <i>Agronomic traits</i> | | |
| EFOSi | <i>Enzyme digestible organic matter at ileum</i> | | |
| EFOSsvin | <i>Enzyme digestible organic matter in pigs</i> | | |
| Erucasyre | <i>Erucic acid</i> | | |
| FEso pr. hkg | <i>Feed units, adult pigs</i> | | |
| FEsv pr. hkg | <i>Feed units, growing pigs</i> | | |
| fht | <i>Index</i> | | |
| Faldtal | <i>Falling number</i> | | |
| Foderkvalitet | <i>Feed quality</i> | | |
| Frøkvalitet | <i>Seed quality</i> | | |
| Frøvægt | <i>Seed weight</i> | | |
| Glucosinolatindhold | <i>Glucosinolate content</i> | | |
| Gluten i kerner (14% vand) | <i>Gluten content in grains at 14 % water</i> | | |
| Gns. | <i>Average</i> | | |
| Gråplet/brunplet | <i>Septoria tritici/Stagonospora nodorum</i> | | |
| Gulrust | <i>Yellow rust (Puccinia striiformis)</i> | | |
| hkg/ha korrigeret til 85 % tørstof | <i>hkg/ha adjusted to 85% dry matter</i> | | |
| Hvedebladplet | <i>Tan spot, DTR (Pyrenophora tritici-repentis)</i> | | |
| Karakter | <i>Score</i> | | |
| Kerne­kvalitet | <i>Grain quality</i> | | |
| Klæbrighed | <i>Stickyness</i> | | |
| Kornvægt, mg pr. korn | <i>Thousand kernel weight (mg/kg)</i> | | |
| Kvalitetsegenskaber | <i>Quality traits</i> | | |
| Led | <i>Entry</i> | | |
| Lejesæd | <i>Lodging</i> | | |
| Linolénsyre | <i>Linolenic acid</i> | | |
| Linolsyre | <i>Linoleic acid</i> | | |
| Meldug | <i>Mildew (Erysiphe graminis)</i> | | |
| Meludbytte | <i>Flour yield</i> | | |
| Modning, dato | <i>Ripeningdate</i> | | |
| Nedknækning, aks | <i>Necking</i> | | |
| Nedknækning, strå | <i>Brackling</i> | | |
| Olieindhold | <i>Oil content</i> | | |
| Oliesyre | <i>Oleic acid</i> | | |
| Plantehøjde | <i>Plant height</i> | | |
| Proteinindhold, pct. | <i>Protein content</i> | | |
| Ramularia | <i>Ramularia (Ramularia collo-cygni)</i> | | |
| Rumvægt, g pr. liter | <i>Specific weight</i> | | |
| Sedimentation | <i>Zeleny sedimentation value</i> | | |
| Skala | <i>Scale</i> | | |
| Skoldplet | <i>Leaf Blotch (Rhynchosporium secalis)</i> | | |
| Sort | <i>Variety</i> | | |
| Sort., pct. kerner>2,5 mm | <i>Grading, pct. kernels > 2.5 mm</i> | | |
| Sort., pct. kerner>2,8 mm | <i>Grading, pct. kernels > 2.8 mm</i> | | |
| Sortskode | <i>Variety code</i> | | |
| Stabilitet | <i>Stability</i> | | |
| Standardkvalitet | <i>Standard quality</i> | | |